

# A revisionary study of *Trachymyrmex zeteki* (Formicidae: Attini) and description of a new species, *Trachymyrmex fovater* sp. n.

Cody R. Cardenas<sup>1</sup>, Alexandria DeMILto<sup>1</sup>, Ted R. Schultz<sup>2</sup>, Rachelle M. M. Adams<sup>1,2</sup>

<sup>1</sup> The Ohio State University, Columbus, Ohio; <sup>2</sup> Smithsonian Natural History Museum, Washington, D. C.

**Introduction** Fungus-growing ants are ubiquitous Neotropical insects with impressive diversity and fascinating evolutionary history. For 65 million years, ca. 250 species of fungus-growing ants have been refining their agricultural practices. *Trachymyrmex zeteki* Weber 1940 has been confused with a sister species due to confounding species descriptions, nearly identical morphological characteristics and overlapping habitats. We propose a new species, *T. fovater* sp. n., to resolve this taxonomic conundrum.

## Statistical Analysis




Figure 1 – Nine standard measurements and five extrapolated indices were used to differentiate species for all castes. Two measurements are included above (n= 106 *T. fovater*, 45 *T. zeteki*). X indicates the measurement of the *T. zeteki* type. A) Total length (Z-test; P< 0.001) B) Weber's length (Z-test; P< 0.001)

## Molecular Analysis





Figure 2 – Two distinct clades of *T. fovater* sp. n. and *T. zeteki* emerged from our phylogenetic analyses of 15 colonies (397 bp of COI).

## Selected Morphological Differences



### *Trachymyrmex zeteki*

- A - Simple setae present mesad of the frontal carinae
- B - Eyes' anterior half surpasses the lateral margin of the head
- C - Distinct acumination on the anterior-lateral margin of the frontal lobes
- D - Antennal scapes, when lodged in antennal scrobe, reach, but do not surpass occipital corners
- E - Deep and wide vertex impression

### *Trachymyrmex fovater* sp. n.

- A - Spatulate bi-colored setae mesad of the frontal carinae
- B - More than half of the eyes' anterior half surpasses the lateral margin of the head
- C - Weak acumination on the anterior-lateral margin of the frontal lobe
- D - Antennal scapes, when lodged in antennal scrobe, surpass occipital corners of the head
- E - Shallow and narrow vertex impression




Figure 3 – Morphology of the worker head capsule referenced. Table 1 & Figure 4 – Illustrates key diagnostic characters which are referenced in Table 1 for comparison to Figure 4.

## Methods




Figure 5 – Morphology: Nine standard measurements  
 Molecular: Extracted DNA from 40 colonies using a Qiagen Blood and Tissue kit. COI gene was amplified using Jerry/Ben Primers (Simon et al. 1994). OSU's Plant-Microbe genomics facility was used for sequencing and data analyzed with Geneious R9

## Conclusion

- Based on 32 morphological differences in workers, gynes, and males, we conclude that *Trachymyrmex fovater* sp. n. has been incorrectly identified as *T. zeteki*
- Molecular differences corroborate our morphological evidence and distinguish two clades (Fig. 2)

## Future Directions

- Use additional primers of COI and genes to improve the phylogenetic resolution
- Examine male genitalia to find informative characters that separate the *T. zeteki* clade
- Identify and list all publications that have incorrectly named *T. zeteki*

**Acknowledgements** Funding was provided by the OSU URO Summer Fellowship Research Grant. We thank Konstantinos Giampoudakis for collections during the Tropical Behavioral, Ecology, and Evolution 2011 course (University of Copenhagen), David Culver and Joan M. Herbers for microscopes, Steve Passoa for technical microscope assistance, the Daly Lab for lab materials, and the Triplehorn Insect Collection for curation materials and use of auto montage equipment.